

Evaluation of Reversed Administration Order of Busulfan (BU) and Cyclophosphamide (CY) as Conditioning on Liver Toxicity in Allogeneic Hematopoietic Stem Cell Transplantation (ALL-HSCT)

Mani Ramzi¹, Nasrin Namdari², Shirin Haghghat³, Hourvash Haghghinejad⁴

¹Hematology and Bone Marrow Transplant Research Center, Shiraz University of Medical Sciences, Shiraz, Iran

²Department of Hematology and Medical Oncology, Shiraz University of Medical Sciences, Shiraz, Iran

³Department of Hematology, Hematology Research Center, Medical Oncology and Stem Cell Transplantation, Shiraz University of Medical Sciences, Shiraz, Iran

⁴Department of Family Medicine, Shiraz University of Medical Sciences, Shiraz, Iran

Corresponding Author: Mani Ramzi, Hematology and Bone Marrow Transplant Research Center, Shiraz University of Medical Sciences, Shiraz, Iran
Email: ramzim@sums.ac.ir

Received: 29, Jun, 2019
Accepted: 15, Feb, 2020

ABSTRACT

Background: Busulfan (BU) in combination with cyclophosphamide (CY) is used as an effective conditioning regimen in hematopoietic SCT. Busulfan, depletes glutathione level in liver and causes elevated levels of CY metabolites. Cyclophosphamide metabolites are highly toxic for sinusoidal endothelial cells and cause VOD/ SOS with high mortality rate.

Materials and Methods: All adult patients with acute leukemia, who were candidates for myeloablative allogeneic SCT admitted in Stem Cell Transplantation center of our center were enrolled in this prospective randomized clinical trial during 2 years. We tested the hypothesis that reverse administration from BU-CY (28 patients) to CY-BU group (27 patients) would reduce liver toxicity.

Results: Liver function tests were significantly higher in BU-CY group between day -1 and +4 ($p < 0.05$), but we do not have VOD/SOS in both groups. The incidence and severity of acute GVHD was higher in BU-CY group, but not statistically significant. Engraftment and mortality rate were not different.

Conclusion: These data support the concept that CY-BU is associated with less liver toxicity, suggesting CY-BU is superior to BU-CY as conditioning.

Keywords: Allogeneic transplant; Conditioning; Busulfan; Cyclophosphamide

INTRODUCTION

Hematopoietic Stem cell Transplantation is a curative therapy for a number of malignant and non-malignant disorders. Busulfan followed by Cyclophosphamide (BU-CY) is used as an effective conditioning regimen in allogeneic hematopoietic Cell Transplantation¹. Liver toxicity and hepatic sinusoidal obstruction syndrome (SOS), also known as Hepatic Veno-occlusive disease (VOD) is a

potentially life-threatening complication that can occur after myeloablative hematopoietic stem cell transplantation²⁻⁵. Incidence of VOD/SOS ranges from 8-14%, and its severe form is associated with mortality rate higher than 80%⁶.

VOD/SOS begins with injury to sinusoidal endothelial cells and hepatocytes due to toxic metabolites generated by high dose alkylating chemotherapy conditioning regimens, such as busulfan,

cyclophosphamide, melphalan, 6-mercaptopurine and possibly Thiotepa.⁷ Busulfan and Cyclophosphamide are metabolized in liver^{8,9}. Busulfan is not toxic for hepatocytes and sinusoidal endothelial cells, but metabolites of cyclophosphamide are highly toxic to sinusoidal endothelial cells¹⁰⁻¹².

Cytotoxic effect of cyclophosphamide is mediated by its active metabolites such as 4-hydroxy cyclophosphamide (4-OHCP) and phosphoramide mustard (PM). The extent of cyclophosphamide metabolism depends, in part, on the activity and concentration of aldehyde dehydrogenase, glutathione and glutathione S-Transferase⁽¹³⁻¹⁵⁾. Glutathione is involved in busulfan clearance, and starting conditioning regimens with busulfan, depletes glutathione level in liver and causes elevated levels of cyclophosphamide metabolites^{11-12,17}.

Restoration of hepatic and sinusoidal endothelial cell glutathione levels prevents injury to hepatic sinusoidal in several different animal models of toxic liver injury¹⁸. Cyclophosphamide administration in less than 24 hours after the last dose of busulfan, causes more mucositis and VOD/SOS because of negative effect on cyclophosphamide pharmacokinetics¹⁹. Studies have suggested less hepatotoxicity when busulfan is administered after cyclophosphamide^{8,19-21}.

Here, we report a prospective randomized clinical trial designed to test the hypothesis that reverse administration of CY and BU reduces hepatotoxicity in comparison to standard BU-CY conditioning regimen.

MATERIALS AND METHODS

All adult patients with acute leukemia, who were candidate for myeloablative allogeneic SCT admitted in Stem Cell Transplantation center of Shiraz were enrolled in this prospective randomized clinical trial during 2 years. They referred to our hospital from September 2013 to September 2015. All eligible patients were provided written informed consent. All procedures were in accordance with the Helsinki protocol of 1975 and approved by Ethics Committee of Shiraz University of Medical Sciences (Shiraz, Iran) Inclusion criteria:

Eligibility criteria included: 1) patients with acute leukemia (ALL and AML) that needs Allogeneic SCT 2) age ≥ 15 years 3) Karnofski performance status of $>70\%$ at the time of HCT

Exclusion criteria:

1) HIV infection 2) chronic liver disease 3) active hepatitis

Conditioning regimen:

Patients randomized to two groups: CY-BU and BU-CY.

Patients in CY-BU group (n=27) received cyclophosphamide on days -8 through -6, and busulfan on days -5 through -2. Transplantation was done on day 0. Patients in control group (BU-CY, n=28) received busulfan on days -8 through -5 and cyclophosphamide on days -4 through -2.

Cyclophosphamide was administered with total dose of 120 mg/kg i.v and busulfan i.v with dose of 0.8 mg/kg/dose every 6 hours, for a total 16 doses. Prophylactic phenytoin was given from days -8 through -2 to all patients. Graft-versus-Host-disease prophylaxis consisted of cyclosporine and methotrexate. Cyclosporin was given from day -1 as an i.v route with 3mg/kg/d, with adjusted doses according to cyclosporine level. Cyclosporin was changed to oral route when the patients could tolerate oral dose. Methotrexate was given at a dose of 10 mg/m² i.v on days +1, +3, +6 and +11. All patients received antifungal, antiviral, antibacterial and ursodiol prophylaxis in accordance to standard practice.

We checked liver function tests on admission, before starting conditioning regimen, one day before transplantation, on transplantation day and on days +4, +8, +12, +16, +20 and +30. The patients were followed till days +100 for GVHD presentation and mortality. Neutrophil engraftment was defined first day of two consecutive days of PMN ≥ 500 /dl.

Statistical analyses

Statistical comparisons of engraftment between two groups was done by independent t test. Frequency of GVHD was compared by chi square test between two groups. Comparison of the liver function tests between two groups was done using Mann-whitney test.

RESULTS

Patients and their characteristics are summarized in Table 1. The median age of patients in BU-CY group was 30 years (range, 15-61 years). Sixteen patients (57.1%) were male and 12(42.9%) were female. Thirteen patients (46.43%) had ALL, 13(46.43%) had AML and 2(7.14%) had MDS. The median age of patients in CY-BU group was 31 years (range: 20-53 years). Seventeen patients (63%) were male and ten (37%) were female. nine (33.3%) had ALL, 18(66.7%) had AML and none of them had MDS.

There was no statistically significant difference between two groups according to age, sex and underlying disease distribution. All transplants were from HLA-matched sibling donors, except one of them, was HLA-matched from his child.

Engraftment

Median times to myeloid engraftment (BU-CY cohort 13 days (range: 9-17) vs CY-BU cohort 13 days (range: 10-18)) were similar between two groups (P-value: 0.93).

Graft Versus Host Disease

The cumulative incidence of acute GVHD at day +100 was 46.6% in the BU-CY group and 37% in CY-BU group (p-value= 0.48). Severity of acute GVHD in BU-CY group was 30.8% grade 1, 38.5% grade 2, 7.7% grade 3, and 23.1% grade 4. None of the patients in CY-BU group had grade 1 and 4 GVHD, 90% had grade 2 and 10% grade 3.

Skin was the most frequent involved organ between two groups. GVHD was more severe in BU-CY group (38.4% grade 3 and 4) compared with CY-BU group (10% grade 3 and 4).

Liver toxicity

Diagnosis of VOD /SOS was not made in both groups. Liver function tests including Alkaline phosphatase and bilirubin were not statistically significant different between two groups, but mean values of ALT on one day before (p=0.02), transplantation day (p=0.025) and day +4 post-transplant (p=0.03) were significantly higher in control group (figure 1). Mean values of AST on one day before (p=.001) and transplantation day (p=0.001) were significantly higher in control group (Figure 1).

Mortality

The cumulative incidence of TRM at day +100 was 10.7% in BU-CY group and 3.7% in CY-BU group (p=0.32). Causes of TRM in the BU-CY group was relapse (1 patient, 3.57%), GVHD (1 patient, 3.57%) and GVHD simultaneous with CMV infection (1 patient, 3.57%), and those in CY-BU group was GVHD (1 patient, 3.7%).

Table1. Patients Characteristics

Characteristics	BU-CY	CY-BU	
Total Patients, n%	28	27	
Sex, n%			0.66
Male	16(57.1%)	17(63%)	
Female	12(42.9)	10(37%)	
Age			0.82
Median (range)	30(15-61)	31(20-53)	
Underlying disease, n%			0.8
All	12(44.4%)	9(33.3%)	
Aml	13(48.1%)	18(66.7%)	
MDs	2(7.4%)	0(0%)	
Donor, n%			0.36
Identical Sibling	26(96.3%)	27(100)	
Other family members	1(3.7%)	0(0%)	
Donor Sex, n%			0.25
Female in male	4(14.8%)	3(11.1%)	
Others	23(85.2%)	24(88.9%)	
Disease condition at Transplantation			0.11
First remission	15(55.6%)	20(74.1%)	
Second remission	3(11.1%)	3(11.1%)	
Others	9(14.8%)	4(14.8%)	








Figure 1. Mean serum levels of AST, ALT, T. bilirubin, D.bilirubin and Alkaline phosphatase for the BU-CY and CY-Bu group at day -8 (before starting the conditioning regimen), -1 (one day before transplantation), 0, +4, +8, +12, +16, +20 and +30.



DISCUSSION

One of the possible approaches to minimizing liver toxicity caused by BU-Cy conditioning is reversing the order of administration, giving CY followed by busulfan.

Median time to myeloid engraftment was not different between BU-CY and CY-BU group in our study ($p= 0.93$). Our results were comparable with those of Cantoni et al. and Kerbauy et al. Myeloid engraftment was similar in mice treated with BU-CY v/s CY-BU conditioning regimen^{8,20}. It seems that reverse administration of conditioning regimen does not affect myeloid engraftment.

The incidence of acute GVHD were similar between two groups (46.4% in BU-CY and 37% in CY-BU, $p=0.48$) in our study, but patients in BU-CY group had more severe GVHD (grade 3 and 4) relative to CY-BU group (38.4% vs 10%). Cantoni et al. demonstrated that patients in BU-CY group had more acute GVHD relative to patients in CY-BU group (75% vs 44%, $p=0.001$), but severity of acute GVHD were similar between two groups⁸.

Liver toxicity and VOD/SOS is a potentially life-threatening complication after myeloablative hematopoietic stem cell transplantation. In our study, mean values of AST and ALT were significantly higher in BU-CY group in comparison to CY-BU group ($p<0.05$). In the study by Cantoni et al. the incidence of SOS was significantly higher in BU-CY group patients (12.5% vs 0% in CY-BU group) ($p= 0.006$). also, serum levels of liver function tests were higher in patients with BU-CY conditioning regimen.⁸In the

study by Kerbauy et al. liver function tests were significantly higher in patients with BU-CY group ($p < 0.05$) and similar results has been reported²⁰.

Rezvani et al. evaluated administering Cyclophosphamide followed by intravenous targeted busulfan in patients with AML, MDS and myelofibrosis. Compared with BU-CY conditioned patients, CY-BU conditioned patients had greater exposure to cyclophosphamide and less exposure to hydroxycyclophosphamide²¹.

In patients with myelofibrosis, incidence of SOS was significantly lower in patients with CY-BU cohort in comparison to BU-CY (0% vs 30%), whereas the incidence of SOS was low in both cohorts with AML and MDS. Rezvani et al. recommends CY-BU as a superior conditioning regimen in patients with myelofibrosis²¹.

CONCLUSION

According to our results and other studies, it seems that CY-BU conditioning is associated with less liver toxicity, VOD/SOS and acute GVHD without interfering with engraftment and mortality rate. Our finding in good concordance of previous studies showed that CY-BU is effective and less toxic and transplant centers can use it as replacement choice of conditioning regime.

ACKNOWLEDGMENT

The authors thank the vice-chancellery of Shiraz University of medical sciences for supporting this research. This article is based on a theses performed by Dr. Nasrin Namdari.

REFERENCES

1. Sadeghi B, Jansson M, Hassan Z, et al. The effect of administration order of BU and CY on engraftment and toxicity in HSCT mouse model. *Bone Marrow Transplant.* 2008; 41(10): 895-904
2. Dalle J, Giralt S. Hepatic Venous-Occlusive Disease after Hematopoietic Stem cell Transplantation: risk factors and Stratification, Prophylaxis, and Treatment. *Biol Blood Marrow Transplant.* 2016; 22(3): 400-409
3. Brodsky R, Topolsky D, Crilley P, et al. Frequency of veno-occlusive disease of the liver in bone marrow transplantation with a modified busulfan/cyclophosphamide preparative regimen. *Am J Clin Oncol.* 1990; 13(3): 221-5

4. Vassal G, Hartmann O, Benhamou E. Busulfan and veno-occlusive disease of the liver. *Ann Intern Med.* 1990; 112(11): 881.
5. Ljungman P, Hassan M, Bekasy AN, et al. High busulfan concentrations are associated with increased transplant-related mortality in allogeneic bone marrow transplant patients. *Bone Marrow Transplant.* 1997; 20(11): 909-13.
6. Coppell JA, Richardson PG, Soiffer R, et al. Hepatic veno-occlusive disease following stem cell transplantation: incidence, clinical course, and outcome. *Biol Blood Marrow Transplant.* 2010; 16(2): 157-168.
7. Ho VT, Revta C, Richardson PG. Hepatic veno-occlusive disease after hematopoietic stem cell transplantation: Update on defibrotide and other current investigational therapies. *Bone Marrow Transplant.* 2008; 41(3): 229-237.
8. Cantoni N, Gerull S, Heim D, et al. Order of application and liver toxicity in patients given BU and Cy containing conditioning regimens for allogeneic hematopoietic SCT. *Bone Marrow Transplant.* 2011; 46: 344-349.
9. Moore MJ. Clinical pharmacokinetics of cyclophosphamide. *Clin Pharmacokinet.* 1991; 20(3): 194-208.
10. Deleve LD. Cellular target of cyclophosphamide toxicity in the murine liver: role of glutathione and site of metabolic activation. *Hepatology.* 1996; 24(4): 830-837.
11. Deleve LD, Wang X. Role of oxidative stress and glutathione in busulfan toxicity in cultured murine hepatocytes. *Pharmacology.* 2000; 60 (3): 143-54.
12. Hassan Z, Hellstrom-lindberg E, Alsadi S, et al. The effect of modulation of glutathione cellular content on busulphan-induced cytotoxicity on hematopoietic cells in vitro and in vivo. *Bone Marrow Transplant.* 2002; 30(3): 141-7.
13. McDonald Gb, Slattery JT, bouvier ME, et al. Cyclophosphamide metabolism, liver toxicity and mortality following hematopoietic stem cell transplantation. *Blood.* 2003; 101(5): 2043-8.
14. Hassan M, Ljungman P, Ringden o, et al. The effect of busulfan on the pharmacokinetics of cyclophosphamide and its 4-hydroxy metabolite: time interval influence on therapeutic efficacy and therapy related toxicity. *Bone marrow Transplant.* 2000; 25(9): 915-924.
15. Fleming RA. An overview of cyclophosphamide and ifosfamide pharmacology. *Pharmacotherapy.* 1997; 17 (5 Pt 2): 146S-154S.
16. Dirven HA, van Ommen B, van Bladeren PJ. Involvement of human glutathione S-transferase isoenzymes in the conjugation of cyclophosphamide metabolites with glutathione. *Cancer Res.* 1994; 54(23): 6215-20.
17. Hassan Z, Hellstrom-Lindberg E, Alsadi s, et al. The effect of modulation of Glutathione cellular content on

busulfan-induced cytotoxicity on hematopoietic cells in vitro and in vivo. *Bone Marrow Transplant.* 2002; 30(3): 141-7.

18. Wang X, KanelGc, Deleve LD. Support of sinusoidal endothelial cell glutathione prevents hepatic veno-occlusive disease in the rat. *Hepatology.* 2000; 31(2): 428-34.

19. Meresse V, Hartmann O, Vassal G, et al. Risk factors for hepatic veno-occlusive disease after high-dose busulfan-containing regimens followed by autologous bone marrow Transplantation: a study in 136 children. *Bone marrow Transplant.* 1992; 10(2): 135-41.

20. Kerbauy FR, Tirapelli B, Akabane H, et al. The effect of administration order of BU and Cy on toxicity in Hematopoietic SCT in humans. *Bone Marrow Transplant.* 2009; 43(11): 883-5.

21. Rezvani AR, McCune JS, Storer BE. Cyclophosphamide followed by Intravenous Targeted Busulfan for Allogenic Hematopoietic Cell Transplantation: Pharmacokinetics and clinical outcomes. *Biol Blood Marrow Transplant.* 2013; 19(7): 1033-39.

22. Nilsson C, Forsman J, Hassan Z, et al. Effect of altering administration order of busulphan and cyclophosphamide on the myeloablative and immunosuppressive properties of the conditioning regimen in mice. *Exp Hematol.* 2005; 33(3): 380–7.