

## Treatment of acute myeloid leukemia (AML): Recent results and new directions

Thomas Büchner

University of Münster, Germany

Acute myeloid leukemia (AML) is a major indication for hematopoietic stem cell transplantation (HSCT). We here describe more in general the present status and new directions in the therapy for AML.

### Status of survival and cure rates

What is realistically the overall complete remission (CR) rate achieved by chemotherapy in usual modifications? This can be calculated from the combined results of the large multicenter randomized trials published since 1980.<sup>(1-32)</sup>

There is only a modest progress in the CR rates when publications until 1990<sup>(5-9,11,30)</sup> are compared with those after 1990<sup>(1-4, 10,12-29, 31, 32)</sup> (Table 1).

**Table- 1; AML Complete Remission Rate in 16 325 Patients (32 Publications)**

Publication Period	1980 - 90	1991 - 2004
Total	63%	65%
Age < 60	69%	72%
Age 60 +	45%	50%

Beyond the CR rate the rate of continuous complete remissions (CCR) at 4-5 years calculated for the numbers of patients who went into CR is relevant to estimate the definite cure rate.

Unlike the CR rate a therapeutic progress over time is seen in the CCR rate almost restricted, however, to the younger age group (Table 2).

**Table- 2; AML Continuous Complete Remissions at 4 - 5 Years (32 Publications)**

Publication Period	1980 - 90	1991 - 2004
Total	16%	27%
Age < 60	17%	32%
Age 60 +	11%	14%

### Role of chemotherapy intensity

The longterm outcome reflected by CCR or relapse-free survival (RFS) appears to depend on the intensity of chemotherapy both in terms of its dosage and its duration, exemplified by 3 sequential trials of the German AML Cooperative group (Figure 1). No consolidation or maintenance in a non-randomized setting is not associated with long-term RFS.<sup>(7)</sup> Consolidation and maintenance (for 3 years) produces superior RFS compared to consolidation alone.<sup>(7)</sup> And the novel strategy of double induction<sup>(18)</sup> followed by consolidation and maintenance is superior to the same sequence with 1 additional course of intensive consolidation instead of maintenance.<sup>(19)</sup>

Notably, the maintenance effect is even seen in patients at high risk according to age 60+, unfavorable karyotype, LDH>700U, and delayed bone marrow blast clearance.<sup>(19)</sup> Thus, a poor prognosis can be improved by intensified chemotherapy. The maintenance effect is even seen in the high risk group of patients at older age. Actually patients 60 years of age or older contribute 2/3 to the entire AML population.<sup>(33)</sup> An improvement in the older age group would therefore favourably influence the overall results in AML.

Since further intensification in myelotoxic chemotherapy is limited, new perspectives lie in more specific, targeted therapies.

## Acute Promyelocytic Leukemia

The best example for a target group gives Acute Promyelocytic Leukemia (APL), pathogenetically associated with the translocation t(15;17) involving the retinoic alpha receptor gene. As shown by all-transretinoic acid (ATRA) the APL blasts are brought to terminal differentiation, where the differentiated granulocytes can still contain multiple Auer rods as they are typical for the immature blasts in APL. Thus, ATRA induces a CR avoiding the stage of bone marrow aplasia. This observation and successful results were first contributed by a group in Shanghai.<sup>(34)</sup> ATRA then formed the basis for its effective combination with cytotoxic agents, mainly idarubicin as in the AIDA (ATRA/idarubicin) regimen by the Italian GIMEMA group<sup>(35)</sup> and similarly by the Spanish PETHEMA group.<sup>(36)</sup> Comparable with these experiences, the combination of ATRA with standard chemotherapy and even high-dose araC produced long-term CCR above 80%.<sup>(37)</sup>

The new competitor to ATRA, Arsenic-Trioxide (ATO) has again been introduced by the group in Shanghai who first used it in APL relapse where they induced high rates of 2<sup>nd</sup> CR.<sup>(38)</sup> The use of ATO in newly diagnosed APL was first investigated by the group in Teheran and proved equivalent to ATRA in the CR rate and duration.<sup>(39)</sup> Meanwhile, the Shanghai group started a randomized trial comparing ATO with ATRA and ATO+ATRA with the latter combination producing 100% CR and no relapse projected to 2 years in limited numbers of patients.<sup>(40)</sup>

## Novel approaches in AML

In Non-APL AML molecularly targeted approaches are right at their beginning and positive outcomes are limited to single cases. Thus, a receptor-tyrosin-kinase-inhibitor targeting c-kit, VEGF and FLT3 induced a sustained remission in a patient with 2<sup>nd</sup> relapse refractory to other options.<sup>(41)</sup>

The immunologically targeted combination of Gemtuzumab/ Ozogamycin (GO) binds to cells with the AML specific CD33 antigen. Given as single agent, GO proved successful in patients with AML relapse<sup>(42)</sup> and patients over 65 years with untreated AML.<sup>(43)</sup>

Targeting multidrug resistance (MDR1) the addition of the MDR inhibitor PSC-833 to chemotherapy did not improve the outcome in patients over 60 years of age (44), while in younger patients of  $\leq 45$  years it appeared to improve the overall and relapse-free survival.<sup>(45)</sup>

In an attempt to enhance the effect of chemotherapy on leukemic cells growth factors were used in several trials mostly not successfully. Recently, the HOVON group demonstrated an improvement of relapse-free survival by G-CSF priming.<sup>(20)</sup> A similar trial by the German AML CG failed to reproduce this effect, so far.<sup>(46)</sup>

## Allogeneic transplantation: new directions

Allogeneic hematopoietic stemcell transplantation (HSCT) can be considered to provide the most potent antileukemic principle through its graft-versus-leukemia (GvL) effect. Unbiased intent-to-treat comparisons of allogeneic HSCT with chemotherapy (figure 2) or autologous HSCT<sup>(47)</sup> actually show significant advantages of allogeneic HSCT in the remission duration whereas advantages in the overall survival failed significance. This shows that allogeneic HSCT is compromised by considerable treatment related mortality (TRM). Attempts to overcome this crucial problem are done by reduced intensity conditioning regimens. Prospective studies using total body irradiation (TBI) are on the way to find the optimal dosage between the conventional 12 Gy and the minimum administered 2 Gy. Preliminary data demonstrating adequate donor-type chimerism and low TRM are promising even in high-risk and/or older patients (personal communication J. Kienast et al.).

## Legends to the figures

### Figure 1

Kaplan-Meier plots of relapse-free survival in adult AML patients at all ages, treated in 3 sequential trials by the German AML Cooperative Group. AMLCG 78, non-randomized trial, patients receiving no consolidation and no maintenance (7). AMLCG 81, randomized trial, patients receiving either consolidation alone or consolidation + maintenance (7). AMLCG 86, randomized trial, patients receiving

double induction, consolidation and either maintenance or intensive consolidation instead of maintenance (8).

**Figure 2**

Intention-to-treat analysis of remission duration according to the presence or absence of a donor among siblings of patients with AML, included in the 1986 trial of the German AML Cooperative Group (unpublished data).

## References

1. Hann IM., Stevens RF, Goldstone AH, et al. Randomized comparison of DAT versus ADE as induction chemotherapy in children and younger adults with acute myeloid leukemia. Results of the Medical Research Council's 10<sup>th</sup> AML Trial (MRC AML10). *Blood* 1997;89:2311-8
2. Burnett AK, Goldstone AH, Stevens RM, et al. Randomised comparison of addition of autologous bone-marrow transplantation to intensive chemotherapy of acute myeloid leukaemia in first remission: results of MRC AML10 trial. *The Lancet* 1998;351:700-8
3. Mandelli F, Vegna ML, Avvisati G, et al. A randomized study of the efficacy of postconsolidation therapy in adult acute nonlymphocytic leukemia: a report of the Italian Cooperative Group GIMEMA. *Annals of Hematology* 1992;64:166-72
4. Cassileth PA, Harrington DP, Appelbaum FR, et al. Chemotherapy compared with autologous or allogeneic bone marrow transplantation in the management of acute myeloid leukemia in first remission. *New England Journal of Medicine* 1998;339:1649-56
5. Rai KR, Hollannd JF, Glidewell OJ, et al. Treatment of acute myeloid leukemia: A study by Cancer and Leukemia Group B. *Blood* 1981;58:1203-12
6. Yates J, Glidewell O, Wiernik P, et al. Cytosine arabinoside with daunorubicin or adriamycin for therapy of acute myelocytic leukemia: A CALGB study. *Blood* 1982;60:454-62
7. Büchner T, Urbanitz D, Hiddemann W, et al. Intensified Induction and Consolidation with or without maintenance chemotherapy for acute myeloid leukemia (AML): Two multicenter studies of the German AML Cooperative Group. *Journal of Clinical Oncology* 1985;3:1583-9
8. Rees JKH, Gray RG, Swirsky D, Hayhoe FGJ. Principal results of the Medical Research Council's 8th acute myeloid leukaemia trial. *The Lancet* 1986;332:1236
9. Hayat M, Jehn U, Willemze R, et al. A randomized comparison of maintenance treatment with androgens, immunotherapy, and chemotherapy in adult acute myelogenous leukemia. *Cancer* 1986;58:617-23
10. Zittoun R, Mandelli F, Willemze R, et al. Autologous or allogeneic bone marrow transplantation compared with intensive chemotherapy in acute myelogenous leukemia. *New England Journal of Medicine* 1995;332:217-223
11. Preisler H, Davis RB, Kirshner J, et al. Comparison of three remission induction regimens and two postinduction strategies for the treatment of acute nonlymphocytic leukemia: A Cancer and Leukemia Group B Study. *Blood* 1987;69:1441-9
12. Hansen OP, Pedersen- Bjergaard J, Ellegaard J, et al. Aclarubicin plus cytosine arabinoside versus daunorubicin plus cytosine arabinoside in previously untreated patients with acute myeloid leukemia: a Danish National Phase III Trial. *Leukemia* 1991;5:510-6
13. Dillman RO, Davis RB, Green MR, et al. A comparative study of two different doses of cytarabine for acute myeloid leukemia: A phase III trial of Cancer and Leukemia Group B. *Blood* 1991;78:2520-6

14. Cassileth PA, Lynch E, Hines JD, et al. Varying intensity of postremission therapy in acute myeloid leukemia. *Blood* 1992;79:1924-30
15. Mayer RJ, Davis RB, Schiffer CA, et al. Intensive postremission chemotherapy in adults with acute myeloid leukemia. *New England Journal of Medicine* 1994;6:896-942
16. Bishop JF, Matthews JP, Young GA, et al. A randomized study of high-dose cytarabine in induction in acute myeloid leukemia. *Blood* 1996;87:1710-7
17. Weick JK, Kopecky KJ, Appelbaum FR, et al. A randomized investigation of high-dose versus standard dose cytosine arabinoside with daunorubicin in patients with previously untreated acute myeloid leukemia: A Southwest Oncology Study Group. *Blood* 1996;88:2841-51
18. Büchner T, Hiddemann W, Wörmann B, et al. Double induction strategy for acute myeloid leukemia: The effect of high-dose cytarabine with mitoxantrone instead of standard- dose cytarabine with daunorubicin and 6-thioguanine. A randomized trial by the German AML Cooperative Group. *Blood* 1999;93:4116-24
19. Büchner T, Hiddemann W, Berdel W, et al. 6-Thioguanine, cytarabine, and daunorubicin (TAD) and high-dose cytarabine and mitoxantrone (HAM) for induction, TAD for consolidation, and either prolonged maintenance by reduced monthly TAD or TAD-HAM-TAD and one course of intensive consolidation by sequential HAM in adult patients at all ages with de- novo acute myeloid leukemia (AML): A randomized trial by the German AML Cooperative Group. *Journal of Clinical Oncology* 2003;21:4496-4504
20. Löwenberg B, van Putten W, Theobald M, et al. Effect of priming with granulocyte colony-stimulating factor on the outcome of chemotherapy for acute myeloid leukemia. *New England Journal of Medicine* 2003;349:743-752
21. Rowe JM, Andersen JW, Mazza JJ, et al. A randomized placebo- controlled phase III study of granulocyte-macrophage colony- stimulating factor in adult patients (>55 to 70 years of age) with acute myelogenous leukemia: A study of the Eastern Cooperative Oncology Group (E1490). *Blood* 1995;86:457-62
22. Witz F, Sadoun A, Perrin MC, et al. A placebo- controlled study of recombinant human granulocyte-macrophage colony- stimulating factor administered during and after induction treatment for de novo acute myelogenous leukemia in elderly patients. *Blood* 1998;91:2722-30
23. Goldstone AH, Burnett AK, Wheatley K, et al. Attempts to improve treatment outcomes in acute myeloid leukemia (AML) in older patients: the results of the United Kingdom Medical Research Council AML 11 trial. *Blood* 2001;98:1302-11
24. Anderson J, Kopecky K, Willmann C, et al. Outcome after induction chemotherapy for older patients with acute myeloid leukemia is not improved with mitoxantrone and etoposide compared to cytarabine and daunorubicin: A Southwest Oncology Group study. *Blood* 2002;100:3869-3876
25. Stone RM, Berg TB, George SL, et al. Granulocyte-macrophage colony-stimulating factor after initial chemotherapy for elderly patients with primary acute myelogenous leukemia. *New England Journal of Medicine* 1995;332:1671-7

26. Stone RM, Berg DT, George SL, et al. Postremission therapy in older patients with de novo acute myeloid leukemia: a randomized trial comparing mitoxantrone and intermediate-dose cytarabine with standard-dose cytarabine. *Blood* 2001;98:548-53
27. Büchner T, Hiddemann W, Wörmann B, et al. Daunorubicin 60 instead of 30 mg/sqm improves response and survival in elderly patients with AML. *Blood* 1997;90 (Suppl 1):583a
28. Dombret H, Chastang C, Fenaux P, et al. A controlled study of recombinant human granulocyte colony-stimulating factor in elderly patients after treatment for acute myelogenous leukemia. *New England Journal of Medicine* 1995;332:1678-83
29. Rees JKH, Gray RG, Weathley K. Dose intensification in acute myeloid leukaemia: greater effectiveness at lower cost. Principal report of the Medical Research Council's AML9 study. *British Journal of Haematology* 1996;94:89-98
30. Bishop JF, Lowenthal RM, Joshua D, et al. Etoposide in acute nonlymphocytic leukemia. *Blood* 1990;75:27-32
31. Löwenberg B, Suci S, Archimbaud E, et al. Mitoxantrone versus daunorubicin in induction-consolidation chemotherapy – The value of low-dose cytarabine for maintenance of remission, and an assessment of prognostic factors in acute myeloid leukemia in the elderly: Final report of the Leukemia Cooperative Group of the European Organisation for the Research and Treatment of Cancer and the Dutch-Belgian Hemato-Oncology Cooperative Hovon Group randomized phase III study AML-9. *Journal of Clinical Oncology* 1998;16:872-81
32. Rowe J, Neuberg D, Friedenber W, et al. A phase 3 study of three induction regimens and of priming with GM-CSF in older adults with acute myeloid leukemia: a trial by the Eastern Cooperative Oncology Group. *Blood* 2004;103:479-485
33. Büchner T, Berdel WE, Wörmann B, et al. Treatment of older patients with AML. *Critical Reviews in Oncology/ Hematology* 2005 (in press)
34. Huang M, Ye Y, Chen S, et al. Use of all- trans retinoic acid in the treatment of acute promyelocytic leukemia. *Blood* 1988; 72:567-572
35. Avvisati G, Lo Coco F, Diverio D, et al. AIDA ( All-trans Retinoic Acid + Idarubicin) in newly diagnosed acute promyelocytic leukemia: a Gruppo Italiano Malattie Ematologiche Maligne Dell' Adulto (GIMEMA) pilot study. *Blood* 1996;88:1390-1398
36. Sanz MA, Martin G, Rayon C, et al. A modified AIDA protocol with anthracycline- based consolidation results in high antileukemic efficacy and reduced toxicity in newly diagnosed PML/RAR  $\alpha$ -positive acute promyelocytic leukemia. *Blood* 1999;94:3015-3021
37. Lengfelder E, Reichert A, Schoch C, et al. Double induction strategy including high dose cytarabine in combination with all- trans retinoic acid: effects in patients with newly diagnosed acute promyelocytic leukemia. *Leukemia* 2000; 14:1362-1370
38. Shen ZX, Chen GQ, Ni JH, et al. Use of arsenic trioxide (As<sub>2</sub>O<sub>3</sub>) in the treatment of acute promyelocytic leukemia (APL). II Clinical efficacy and pharmacokinetics in relapsed patients. *Blood* 1992;79:543-553

39. Gavamzadeh A, Alimoghaddam, K, Aghdami, N, Treatment of acute promyelocytic leukemia by arsenic trioxide. *Proc. ASCO* 2003;22:568
40. Shen ZX, Shi ZZ, Fang J, All- trans retinoic acid/As<sub>2</sub>O<sub>3</sub> combination yields a high quality remission and survival in newly diagnosed acute promyelocytic leukemia. *PNAS* 2004;101:5328-5335
41. Mesters R, Padró T, Bieker R, et al. Stable remission after administration of the receptor tyrosine kinase inhibitor SU5416 in a patient with refractory acute myeloid leukemia. *Blood* 2001;98:241-243
42. Sievers EL, Larson RA, Stadtmauer EA, et al. Efficacy and safety of Gemtuzumab Ozogamicin in patients with CD33-positive acute myeloid leukemia in first relapse. *Journal of Clinical Oncology* 2001;19:3244-3254
43. Nabhan C, Rundhaugen LM, Riley MB, et al. Phase III pilot trial of gemtuzumab ozogamicin (GO) as first line therapy in acute myeloid leukemia patients age 65 or older. *Leukemia Research* 2005; 29:53-57
44. Baer MR, George SL, Dodge RK, et al. Phase 3 study of the multidrug resistance modulator PSC-833 in previously untreated patients 60 years of age and older with acute myeloid leukemia: Cancer and Leukemia Group B study 9720. *Blood* 2002;100:1224-1232
45. Kolitz JE, George SL, Dodge RK, et al. Dose escalation studies of cytarabine, daunorubicin, and etoposide with and without multidrug resistance modulation with PSC-833 in untreated adults with acute myeloid leukemia younger than 60 years: final induction results of cancer and leukemia Group B study 9621. *Journal of Clinical Oncology* 2004;22:4290-4301
46. Büchner T, Berdel WE, Hiddemann W. Priming with granulocyte colony stimulating factor- relation to high- dose cytarabine in acute myeloid leukemia. *New England Journal of Medicine* 2004;350:2215-16 (letter)
47. Suci S, Mandelli F, de Witte T, et al. Allogeneic compared with autologous stem cell transplantation in the treatment of patients younger than 46 years with acute myeloid leukemia (AML) in first complete remission (CR1): an intention-to- treat analysis of the EORTC/GIMEMA AML-10 trial. *Blood* 2003;102:1232-1240